

Phenotypic anchoring of gene expression
changes during estrogen-induced
uterine growth.

Jonathan G. Moggs, Helen Tinwell, Tracey Spurway,
Hur-Song Chang, Ian Pate, Fei Ling Lim,
David J. Moore, Anthony Soames, Ruth Stuckey,
Richard Currie, Tong Zhu, Ian Kimber, John Ashby and
George Orphanides

doi:10.1289/txg.7345 (available at <http://dx.doi.org/>)
Online 7 October 2004



The National Institute of Environmental Health Sciences
National Institutes of Health
U.S. Department of Health and Human Services

Toxicogenomics

Phenotypic anchoring of gene expression changes during estrogen-induced uterine growth.

Jonathan G. Moggs¹, Helen Tinwell¹, Tracey Spurway¹, Hur-Song Chang^{2,3}, Ian Pate¹, Fei Ling Lim¹, David J. Moore¹, Anthony Soames¹, Ruth Stuckey¹, Richard Currie¹, Tong Zhu², Ian Kimber¹, John Ashby¹ and George Orphanides¹.

¹Syngenta Central Toxicology Laboratory, Alderley Park, Cheshire, SK10 4TJ, UK.

²Syngenta Biotechnology Inc., Research Triangle Park, NC 27709, USA. ³Present address: Diversa Corporation, 4955 Directors Place, San Diego, CA 92121, USA.

Correspondence to: G. Orphanides, Syngenta CTL, Alderley Park, Cheshire, SK10 4TJ, UK.

Tel: 441625510803.

Fax: 441625585715.

E-mail: george.orphanides@syngenta.com

Running Title:

uterotrophic transcriptional program

Key words:

estrogen, uterus, microarray, gene expression, phenotypic anchoring

Acknowledgements:

This work was partially supported by the UK Food Standards Agency (FSA). We thank M.G. Parker, D.G. Deavall, N. Wallis, and T. Barlow for critical comments on the manuscript, P. Lefevre and J. Odum for technical assistance and Ilya Kupershmidt and Ewan Hunter (Silicon Genetics) for advice on statistical analysis of microarray data. No conflict of interest was reported.

Abbreviations:

ER – estrogen receptor

ANOVA – analysis of variance

AO – arachis oil

E₂ – 17β-estradiol

sc – subcutaneous

Outline of section headers:

Abstract

Introduction

Materials and Methods

Results

Discussion

References

Tables

Figure Legends

Figures

Abstract

A major challenge in the emerging field of toxicogenomics is to define the relationships between chemically induced changes in gene expression and alterations in conventional toxicological parameters, such as clinical chemistry and histopathology. We have explored these relationships in detail using the rodent uterotrophic assay as a model system. Gene expression levels, uterine weights and histological parameters were analyzed 1, 2, 4, 8, 24, 48, and 72 hr after exposure to the reference physiological estrogen, 17 β -estradiol (E₂). A multi-step analysis method, involving unsupervised hierarchical clustering followed by supervised gene ontology driven clustering, was used to define the transcriptional program associated with E₂-induced uterine growth and to identify groups of genes that may drive specific histologic changes in the uterus. This revealed that uterine growth and maturation is preceded and accompanied by a complex, multi-stage molecular program. The program begins with the induction of genes involved in transcriptional regulation and signal transduction, and is followed, sequentially, by the regulation of genes involved in protein biosynthesis, cell proliferation and epithelial cell differentiation. Furthermore, we have identified genes with common molecular functions that may drive fluid uptake, coordinated cell division and remodeling of luminal epithelial cells. These data define the mechanism by which an estrogen induces organ growth and tissue maturation, and demonstrate that comparison of temporal changes in gene expression and conventional toxicology endpoints can facilitate the phenotypic anchoring of toxicogenomic data.